

24 (New) The stabilized pharmaceutical preparation of claim 1 wherein the α -amino acid is glycine and

wherein,

R_1 is a hydrogen atom, a hydroxyl group, a methyl group or an ethyl group;

R_2 is a monovalent group selected from:

a straight or branched alkyl group of 3 - 8 carbon atoms;

a straight or branched alkylene group of 3 - 8 carbon atoms;

a straight or branched alkyl group of 3 - 8 carbon atoms which is mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 3 - 8 carbon atoms;

a cycloalkyl group of 3 - 8 carbon atoms which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group,

a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms, is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups ($-\text{CH}_2-$) is replaced by $-\text{O}-$, $-\text{NH}-$, $-\text{S}-$, $-\text{SO}-$ or $-\text{S}(\text{O})_2-$, and one or two of the unsubstituted methylene groups ($-\text{CH}_2-$) are mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

C' cont.
a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups ($-\text{CH}_2-$) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by $-\text{O}-$, $-\text{NH}-$, $=\text{N}-$, $-\text{S}-$, $-\text{SO}-$ or $-\text{S}(\text{O})_2-$;

Sub
a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups ($-\text{CH}_2-$) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by $-\text{O}-$, $-\text{NH}-$, $=\text{N}-$, $-\text{S}-$, $-\text{SO}-$ or $-\text{S}(\text{O})_2-$, and one or two of the unsubstituted methylene groups ($-\text{CH}_2-$) being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups ($-\text{CH}_2-$) is replaced by $-\text{O}-$, $-\text{NH}-$, $-\text{S}-$, $-\text{SO}-$ or $-\text{S}(\text{O})_2-$;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, said phenyl group being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

C1 cont.
a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, one of the methylene groups

(-CH₂-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, and one of the methylene groups (-CH₂-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)₂- and one or two of the unsubstituted methylene groups (-CH₂-) being mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a phenyl or naphthyl group;

a phenyl group substituted with a methylenedioxy group;

a phenyl or naphthyl group which is mono-, di- or tri-substituted with a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group, a phenoxy group, a phenylmethoxy group, a phenylmethoxy group wherein said phenyl ring is mono-substituted with a halogen atom, trifluoromethyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring, a cycloalkanediylmethoxy group having 5 - 8 carbon atoms in the cycloalkanediyl ring, a cycloalkylmethoxy

group wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, a cycloalkenylmethoxy group wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, a cycloalkanediénylmethoxy group wherein one of the methylene groups (-CH₂-) in said cycloalkanediényl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂- group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring wherein said cycloalkyl ring is mono-substituted with a halogen atom, trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH₂-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring wherein said cycloalkenyl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH₂-) in said cycloalkenyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, or a cycloalkanediénylmethoxy group having 5 - 8 carbon atoms in the cycloalkanediényl ring wherein said cycloalkanediényl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a

hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH₂-) in said cycloalkanedieryl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS-;

an -O-, -S- or -SS-phenyl group;

a diphenylamino group;

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an -O-, -S- or -SS-phenyl group wherein said phenyl group is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group,

a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

or

R₁ and R₂, together with the carbon atom to which they are attached, may form a divalent group selected from:

a cycloalkylidene group of 5 - 8 carbon atoms;

a cycloalkylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group or a carboxyl group;

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂- group and one or more of the unsubstituted methylene groups (-CH₂-) in said cycloalkyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

C' cont
Sub
(1)
a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂- group and one or more of the unsubstituted methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms;


a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms, said phenyl ring being mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedierylidene group of 5 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedierylidene group of 5 - 8 carbon atoms, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group.

Please amend claim 1 as follows.

1. (Twice Amended) A stabilized pharmaceutical preparation, comprising: (a) and α amino acid; (b) an optional auxiliary agent for manufacturing a pharmaceutical preparation; and (c) a 4-amino-3-substituted-butanoic acid derivative, which 4-amino-3-substituted-butanoic acid derivative has the general formula:



wherein,

R₁ is a hydrogen atom, a hydroxyl group, a methyl group or an ethyl group;

R₂ is a monovalent group selected from:

a straight or branched alkyl group of 3 - 8 carbon atoms;

a straight or branched alkylene group of 3 - 8 carbon atoms;

a straight or branched alkyl group of 3 - 8 carbon atoms which is mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 3 - 8 carbon atoms;

a cycloalkyl group of 3 - 8 carbon atoms which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group,

a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms, is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups ($-\text{CH}_2-$) is replaced by $-\text{O}-$, $-\text{NH}-$, $-\text{S}-$, $-\text{SO}-$ or $-\text{S}(\text{O})_2-$, and one or two of the unsubstituted methylene groups ($-\text{CH}_2-$) are mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups ($-\text{CH}_2-$) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by $-\text{O}-$, $-\text{NH}-$, $=\text{N}-$, $-\text{S}-$, $-\text{SO}-$ or $-\text{S}(\text{O})_2-$;

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups ($-\text{CH}_2-$) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by $-\text{O}-$, $-\text{NH}-$, $=\text{N}-$, $-\text{S}-$, $-\text{SO}-$ or $-\text{S}(\text{O})_2-$, and one or two of the unsubstituted methylene groups ($-\text{CH}_2-$) being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups ($-\text{CH}_2-$) is replaced by $-\text{O}-$, $-\text{NH}-$, $-\text{S}-$, $-\text{SO}-$ or $-\text{S}(\text{O})_2-$;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, said phenyl group being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

C2 cont
Sub
(1)
a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, one of the methylene groups

(-CH₂-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, and one of the methylene groups (-CH₂-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)₂- and one or two of the unsubstituted methylene groups (-CH₂-) being mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a phenyl or naphthyl group;

a phenyl group substituted with a methylenedioxy group;

a phenyl or naphthyl group which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group, a phenoxy group, a phenylmethoxy group, a phenylmethoxy group wherein said phenyl ring is mono-substituted with a halogen atom, trifluoromethyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring, a cycloalkanediethylmethoxy group having 5 - 8 carbon atoms in the cycloalkanediethyl ring, a

cycloalkylmethoxy group wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, a cycloalkenylmethoxy group wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, a cycloalkanedieryl-methoxy group wherein one of the methylene groups (-CH₂-) in said cycloalkanedieryl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂- group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring wherein said cycloalkyl ring is mono-substituted with a halogen atom, trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH₂-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring wherein said cycloalkenyl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH₂-) in said cycloalkenyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, or a cycloalkanedierylmethoxy group having 5 - 8 carbon atoms in the cycloalkanedieryl ring wherein said cycloalkanedieryl ring is

02 cont.
SUB
11

mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH₂-) in said cycloalkanedieryl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS-;

an -O-, -S- or -SS-phenyl group;

a diphenylamino group;

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, a alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an -O-, -S- or -SS-phenyl group wherein said phenyl group is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

or

R₁ and R₂, together with the carbon atom to which they are attached, may form a divalent group selected from:

a cycloalkylidene group of 5 - 8 carbon atoms;

a cycloalkylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group or a carboxyl group;

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂- group and one or more of the unsubstituted methylene groups (-CH₂-) in said cycloalkyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino

group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

C² cont
sub
Q1
a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂- group and one or more of the unsubstituted methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms, said phenyl ring being mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

provided that when R_2 is a phenyl or naphthyl group which is mono-, di- or tri-substituted with a halogen atom, the α -amino acid is not glycine; and

provided that when the 4-amino-3-substituted-butanoic acid derivative is gabapentin, the α -amino acid is not methyl-D-aspartic acid.

C₂ cont.

Sub
11/11